




ΕΛΛΗΝΙΚΗ ΡΕΥΜΑΤΟΛΟΓΙΚΗ ΕΤΑΙΡΕΙΑ
& ΕΠΑΓΓΕΛΜΑΤΙΚΗ ΕΝΩΣΗ ΡΕΥΜΑΤΟΛΟΓΩΝ ΕΛΛΑΔΟΣ



Νεότερες εξελίξεις στη Ρευματολογία

19 - 12 - 2022

Αρτηρίτιδα Takayasu (TAK)

2022 AMERICAN COLLEGE OF RHEUMATOLOGY / EULAR CLASSIFICATION CRITERIA FOR TAKAYASU ARTERITIS

CONSIDERATIONS WHEN APPLYING THESE CRITERIA

- These classification criteria should be applied to classify the patient as having Takayasu arteritis when a diagnosis of medium-vessel or large-vessel vasculitis has been made
- Alternate diagnoses mimicking vasculitis should be excluded prior to applying the criteria

ABSOLUTE REQUIREMENTS

Age \leq 60 years at time of diagnosis
Evidence of vasculitis on imaging¹

ADDITIONAL CLINICAL CRITERIA

Female sex	+1
Angina or ischemic cardiac pain	+2
Arm or leg claudication	+2
Vascular bruit ²	+2
Reduced pulse in upper extremity ³	+2
Carotid artery abnormality ⁴	+2
Systolic blood pressure difference in arms \geq 20 mm Hg	+1

ADDITIONAL IMAGING CRITERIA

Number of affected arterial territories (select one) ⁵	
One arterial territory	+1
Two arterial territories	+2
Three or more arterial territories	+3
Symmetric involvement of paired arteries ⁶	+1
Abdominal aorta involvement with renal or mesenteric involvement ⁷	+3

Sum the scores for 10 items, if present. A score of \geq 5 points is needed for the classification of TAKAYASU ARTERITIS.

Σκορ \geq 5 = TAK

Grayson PC, et al. *Arthritis Rheumatol* 2022;74:1872-1880
doi: [10.1002/art.42324](https://doi.org/10.1002/art.42324) (ελεύθερα)

Γιγαντοκυτταρική αρτηρίτιδα (GCA)

2022 AMERICAN COLLEGE OF RHEUMATOLOGY / EULAR CLASSIFICATION CRITERIA FOR GIANT CELL ARTERITIS

CONSIDERATIONS WHEN APPLYING THESE CRITERIA

- These classification criteria should be applied to classify the patient as having giant cell arteritis when a diagnosis of medium-vessel or large-vessel vasculitis has been made
- Alternate diagnoses mimicking vasculitis should be excluded prior to applying the criteria

ABSOLUTE REQUIREMENT

Age \geq 50 years at time of diagnosis

ADDITIONAL CLINICAL CRITERIA

Morning stiffness in shoulders/neck	+2
Sudden visual loss	+3
Jaw or tongue claudication	+2
New temporal headache	+2
Scalp tenderness	+2
Abnormal examination of the temporal artery ¹	+2

LABORATORY, IMAGING, AND BIOPSY CRITERIA

Maximum ESR \geq 50 mm/hour or maximum CRP \geq 10 mg/liter ²	+3
Positive temporal artery biopsy or halo sign on temporal artery ultrasound ³	+5
Bilateral axillary involvement ⁴	+2
FDG-PET activity throughout aorta ⁵	+2

Sum the scores for 10 items, if present. A score of \geq 6 points is needed for the classification of GIANT CELL ARTERITIS.

Σκορ \geq 6 = GCA

Ponte C, et al. *Arthritis Rheumatol* 2022;74:1881-89
doi: [10.1002/art.42325](https://doi.org/10.1002/art.42325) (ελεύθερα)

Σημείωση: Τα παρόντα κριτήρια, ως κριτήρια ταξινόμησης και όχι διάγνωσης, πρέπει να εφαρμόζονται αφού σε έναν ασθενή έχει τεθεί η διάγνωση της αγγειίτιδας μεγάλου ή μεσαίου μεγέθους αγγείων, για την ταξινόμηση σε μία από τις τρεις νόσους. Καταστάσεις που μιμούνται αγγειίτιδα (vasculitis mimics) πρέπει να έχουν αποκλειστεί πριν την εφαρμογή των κριτηρίων.